

Hearing Health Hour Webinar Bibliography

Ménière's Disease | Monday, October 17, 5pm ET

Presenter: Wafaa Kaf, M.D., Ph.D., CCC-A, FAAA

- Al-momani, M., Ferraro, J., Gajewski, B., & Ator, G. (2009) Improved sensitivity of electrocochleography in the diagnosis of Meniere's disease, *International Journal of Audiology*, 48:11, 811-819
- Balaban, C. (2013). A Neurophysiologic Explanation of Illness Behaviors Commonly Seen in Persons with Vestibular Disorders. *Vestibular Rehabilitation: An Advanced Course & Update*. University of Pittsburgh. Volume 1. May 17-19, 2013.
- Basura GJ, Adams ME, Monfared A, et al. (2020) Clinical practice guideline: ménière's disease. *Otolaryngol Head Neck Surg*. 2020;162(2_Suppl):S1-S55.
<https://www.entnet.org/quality-practice/quality-products/clinical-practice-guidelines/menieres-disease/>
- Claes, G. M., Wyndaele, M., De Valck, C. F., Claes, J., Govaerts, P., Wuyts, F. L., & Van de Heyning, P. H. (2008). Travelling wave velocity test and Ménière's disease revisited. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*, 265(5), 517–523. <https://doi.org/10.1007/s00405-007-0486-7>
- Corrales, C.E., & Mudry A. (2017). History of the endolymphatic sac: from anatomy to surgery. *Otol Neurotol* 38:152-156.
- Ferraro JA. Electrocochleography: a review of recording approaches, clinical applications, and new findings in adults and children. *J Am Acad Audiol*. 2010 Mar;21(3):145-52.
- Ferraro, J., Kileny, P., Grasel, S., & Tognola, G. (2019). Electrocochleography: New Uses for an Old Test and Normative Values. *Am J Audiol*, 28(S3)
- Güneri, E. A., Çakır, A., & Mutlu, B. (2016). Validity and Reliability of the Diagnostic Tests for Ménière's Disease. *Turkish archives of otorhinolaryngology*, 54(3), 124–130. <https://doi.org/10.5152/tao.2016.1697>
- Hain, T. (2022). Fluid pathways in the inner ear. *Dizziness-and-balance.com*. Accessed September 10, 2022. <https://dizziness-and-balance.com/anatomy/ear/endolymphatic-sac.htm>
- Hornibrook, J., Coates, M., Goh, A., Gourley, J., & Bird, P. (2012). Magnetic resonance imaging for Ménière's disease: correlation with tone burst electrocochleography. *The Journal of laryngology and otology*, 126(2), 136–141. <https://doi.org/10.1017/S0022215111003112>
- Lefler, S., Kaf, W.A., & Ferraro, J. (2021). Comparing Simultaneous Electrocochleography and Auditory Brainstem Response Measurements using Three Different Extra-Tympanic Electrodes. *J Am Acad Audiol*, 32(6), 339-346.
- Liu, F., Huang, W., Chen, Q., Meng, X., Wang, Z., & He, Y. (2014). Noninvasive evaluation of the effect of endolymphatic sac decompression in Ménière's disease using magnetic resonance imaging. *Acta Oto-Laryngologica*, 134(7), 666–671. <https://doi.org/10.3109/00016489.2014.885118>

- Loureiro, R. M., Sumi, D. V., Lemos, M. D., Tames, H. L. V. C., Gomes, R. L. E., Danie I, M. M., Soares, C. R., Murakoshi, R. W., & Funari, M. B. G. (2019). The role of magnetic resonance imaging in Ménière disease: the current state of endolymphatic hydrops evaluation. *einstein (São Paulo)*, 17(1), eMD4743. https://doi.org/10.31744/einstein_journal/2019MD4743
- Kaf, W.A. (2018). Moving from Bedside to Clinic: Electrocochleography Applications. *Canadian Audiologist*, 5(5). <http://canadianaudiologist.ca/issue/volume-5-issue-5-2018/electrocochleography-feature-2/>
- Kaf, W.A., Lewis, K.M., Yavuz, E., Dixon, S.M., van Ess, M., Jamos, A.M., & Delgado, R.E. (2017). Fast click rate electrocochleography and auditory brainstem response in normal-hearing adults using Continuous Loop Averaging Deconvolution. *Ear Hear*, 38(2), 244-254.
- Kennedy, A., Kaf, W.A., Ferraro, J., Delgado, R., & Lichtenhan, J. (2017). Human summing potential amplitudes vary with tone burst repetition rate and duration. *Frontiers in Neuroscience*, volume 11, articles 429, 1-11. <https://doi.org/10.3389/fnins.2017.00429>
- Megerian C.A., Heddon C., Melki S., Momin S., Paulsey J., Obokhare J., Alagramam K. (2010). Surgical Induction of Endolymphatic Hydrops by Obliteration of the Endolymphatic Duct. *JoVE*. 35. <http://www.jove.com/index/Details.stp?ID=1728>, doi: 10.3791/1728
- Melki, S. J., Li, Y., Semaan, M. T., Zheng, Q. Y., Megerian, C. A., & Alagramam, K. N. (2014). A mouse model validates the utility of electrocochleography in verifying endolymphatic hydrops. *Journal of the Association for Research in Otolaryngology : JARO*, 15(3), 413–421.
- Pyykkö, I., Pyykkö, N., Zou, J., & Manchaiah, V. (2022). Characterization of Balance Problems and Rehabilitation Needs of Patients with Ménière's Disease. *Audiology Research*, 12(1), 22–32. <https://doi.org/10.3390/audiolres12010003>
- Rauch, S.D., Zhou, G., Kujawa, S., Guinan, J., & Herrmann, B. (2004). Vestibular Evoked Myogenic Potentials Show Altered Tuning in Patients with Meniere's Disease. *Otology & Neurotology*, 25: 333–338
- Salt AN, Plontke SK. Endolymphatic hydrops: pathophysiology and experimental models. *Otolaryngol Clin North Am*. 2010 Oct;43(5):971-83. doi: 10.1016/j.otc.2010.05.007. PMID: 20713237; PMCID: PMC2923478.
- Sharon, J. D., Trevino, C., Schubert, M. C., & Carey, J. P. (2015). Treatment of Ménière's Disease. *Current treatment options in neurology*, 17(4), 341. <https://doi.org/10.1007/s11940-015-0341-x>
- Ward, B., Wettstein, V., Golding, J., Corallo, G., Nuti, D., Trabalzini, F., & Mandalà, M. (2019). Patient Perceptions of Effectiveness in Treatments for Meniere's Disease: a National Survey in Italy. *Journal of International Advanced Otology*, 15(1), 112–117. <https://doi.org/10.5152/iao.2019.5758>